

# Impact of horizontal resolution and model time step on European precipitation extremes in the OpenIFS 43r3 atmosphere model

Yingxue Liu<sup>1,2</sup>, Joakim Kjellsson<sup>1,2</sup>, Abhishek Savita<sup>1</sup> and Wonsun Park<sup>3,4</sup>

<sup>1</sup>GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

<sup>2</sup>Faculty of Mathematics and Natural Sciences, Christian Albrechts University of Kiel, Kiel, Germany

<sup>3</sup>Center for Climate Physics, Institute for Basic Science (IBS), Busan, Republic of Korea

<sup>4</sup>Department of Climate System, Pusan National University, Busan, Republic of Korea

*Correspondence to:* Yingxue Liu (yliu@geomar.de)

**Abstract:** Events of extreme precipitation pose a hazard to many parts of Europe but are typically not well represented in climate models. Here, we evaluate daily extreme precipitation over Europe during 1982–2019 in observations (GPCC), reanalysis (ERA5) and a set of atmosphere-only simulations at low- (100 km), medium- (50 km) and high- (25 km) horizontal resolution with identical vertical resolutions using OpenIFS (version 43r3). We find that both OpenIFS simulations and reanalysis underestimate the rates of extreme precipitation compared to observations. The biases are largest for the lowest resolution (100 km) and decrease with increasing horizontal resolution (50 and 25 km) simulations in all seasons. The sensitivity to horizontal resolution is particularly high in mountain regions (such as the Alps, Scandinavia, Iberian Peninsula), likely linked to the sensitivity of vertical velocity to the representation of topography. The sensitivity of precipitation to model resolution increases dramatically with increasing percentiles, with modest biases in the 70<sup>th</sup>–80<sup>th</sup> percentile range and large biases above the 99<sup>th</sup> percentile range. We also find that precipitation above the 99<sup>th</sup> percentile mostly consists of large-scale precipitation (~80 %) in winter, while in summer it is mostly large-scale precipitation in Northern Europe (~70 %) and convective precipitation in Southern Europe (~70 %). Compared to ERA5, the OpenIFS overestimates large-scale precipitation extremes in winter, but underestimates in summer. The discrepancy between OpenIFS and ERA5 decreases with increasing horizontal resolutions. We also examine the sensitivity of extreme precipitation to model time step and find that the convective contribution to extreme precipitation is more sensitive to the model time step than the horizontal resolution. This is likely due to the sensitivity of convective activity to model time step. On the other hand, the large-scale contribution to extreme precipitation is more sensitive to horizontal resolution than the model

time step, which may be due to sharper fronts and steeper topography at higher horizontal resolution.

## 1. Introduction

Extreme precipitation events have severe impacts on our society and ecosystems. For example, extreme precipitation caused a devastating flood in Germany in 2021, in which around 180 people died. The frequency and intensity of extreme precipitation are projected to increase over most regions in the future (Myhre et al., 2019, Intergovernmental Panel on Climate Change, 2023; Li et al., 2021). The increasing extreme precipitation poses a threat for society and must thus be realistically simulated and projected accurately for future climates. However, the climate models have large biases in simulating extreme precipitation events due to coarse horizontal resolution grid and long model time step etc. (Alexander et al., 2019; Avila et al., 2015; Sillmann et al., 2013). The model biases are also hard to evaluate as we lack long-term observations. This study aims to understand the sensitivity of extreme precipitation to model horizontal resolution and model time step.

Extreme precipitation events are usually underestimated in CMIP models (O’Gorman, 2015; Sillmann et al., 2013). Some studies found the simulated extreme precipitation at higher atmosphere horizontal resolutions is more realistic (Wehner et al., 2010, 2014). Jong et al. (2023) found that the characteristics of extreme precipitation at 25 km resolution configurations have smaller biases than at 50 and 100 km. While, Kopparla et al. (2013) found that the reduced extreme precipitation biases at higher horizontal resolution do not hold for all regions (e.g., Australia).

Strandberg and Lind (2021) reported the effect of horizontal resolution on European extreme precipitation is largest in regions with complex topography and in the summer season when precipitation is mostly caused by convective processes using coupled models, in agreement with Iles et al. (2020). However, Li et al. (2011) demonstrated that the impact of horizontal resolution on global precipitation extremes is manifested mostly by its effects on large-scale precipitation, which could be due to the improved large-scale circulation (Hack et al., 2006). Other studies also found an increasing large-scale precipitation with higher resolution but the convective precipitation is rather insensitive to resolution (Jung et al., 2012; Kopparla et al., 2013; Bacmeister et al., 2013).

No global atmosphere-model simulations explicitly resolve all the physical processes and all must therefore employ parametrizations of such motions. Both the individual physical parameterization and their coupling to the dynamics can benefit from a shorter time step (Jung et al., 2012). Mishra & Sahany (2011) found a more realistic simulation of the heavy precipitation in the tropics when the time step was shortened from 60 min to 5 min at a coarse (~300 km) resolution in a short-period (12 months) configuration. Jung et al. (2012) reported that the biases of mean precipitation were smaller at 60 minutes than at 15 minutes with 126 km resolution in IFS model. However, Roberts et al. (2018) found a minimal impact on model biases when shorten the time step from 20 to 15 minutes at 25 km in IFS model. They either did not investigate the multi-year precipitation extremes, or did not explore the extremes in IFS model.

A recent study by Savita et al. (2024) explored the sensitivity of global mean precipitation to the horizontal resolution and model time step in atmosphere-only simulations with OpenIFS. However, the extreme precipitation's sensitivity to horizontal resolution and time step was not investigated. In this study, we investigated the impact of horizontal resolutions (~100 km, ~50 km, and ~25 km) and model time steps (60 minutes, 30 minutes, and 15 minutes) on daily extreme precipitation using OpenIFS simulations and compare them with observation. We also studied on the convective and large-scale precipitation in all simulations. Precipitation extremes sensitivity to model time step is the first time explored in this work using 100 km in OpenIFS atmosphere model. Besides the extremes, we also explored multi-percentile precipitation's sensitivity to horizontal resolution and time step. This paper is structured as follows: section 2 describes the data and methodology, and section 3 discusses the results. The conclusion and discussion can be found in section 4.

## 2. Data and Methods

### 2.1 Model, observation, and reanalysis data

The OpenIFS is derived from the Integrated Forecasting System at the European Centre for Medium-range Weather Forecasting (ECMWF-IFS) cycle 43 release 3 (43r3) (ECMWF, 2017). We use the same AMIP simulations that were used in Savita et al. (2024) which cover the

**Deleted:** Extreme precipitation events have severe impacts on human society and ecosystems. For example, Germany experienced extreme precipitation during mid-July 2021, which exceeded 100 mm/d over a large area resulting in a devastating flood. The recent flood is one of the most serious natural disasters for Germany since 1962, in which around 180 people died as a result of the flood. Coupled Model Intercomparison Project (CMIP) models are used to understand the present and future climate. The CMIP5 models project that the frequency of the most intense precipitation observed today in Europe would be almost double in the future at each 1°C of warming (Myhre et al., 2019). Recently the CMIP6 models also projected an increase in extreme precipitation over most of the regions under global warming (Intergovernmental Panel on Climate Change, 2023; Li et al., 2021). The increasing extreme precipitation poses a threat for society and must thus be realistically simulated and projected accurately for future climates. However, the climate models have large uncertainties in simulating extreme precipitation events due to lack of observations, coarse horizontal resolution grid, long model time step etc. (Alexander et al., 2019; Avila et al., 2015; Sillmann et al., 2013). This study aims to understand the sensitivity of extreme precipitation to model resolution and time step. The CMIP models can simulate time-mean precipitation very well but usually underestimate the intensity and frequency of extreme precipitation (O'Gorman, 2015; Sillmann et al., 2013). The intensity of simulated extreme precipitation often increases with increased horizontal resolution in atmosphere models (Caldwell, 2010; Rauscher et al., 2016; Wehner et al., 2010, 2014). Jong et al. (2023) analyzed the extreme precipitation in Northeastern United States (US) using the Seamless system for Prediction and Earth system Research (SPEAR) and Large Ensemble of Community Earth System Model version 1 (CESM-LE) model simulations at different horizontal resolutions, and they found that a model with 25 km horizontal resolution simulates much more realistic frequency, amplitude, temporal variability and trends in extreme precipitation than 50 and 100 km model resolution. However, Koppala et al. (2013) found that the reduced biases at higher horizontal resolution do not hold for all regions. They concluded that extreme precipitation with finer model resolutions in Community Atmospheric Model version 4 (CAM4) has better agreement with observational datasets in Europe and the US, but not in Australia. Considering the time and computational cost, climate simulations of more than 100 years are generally not feasible with high-resolution (25 km or higher) models. Instead, regional climate models (RCMs) are developed by focusing on a particular region, where higher-resolution model simulations can be conducted with reduced cost (Laprise, 2008). Strandberg & Lind (2021) compared the precipitation using both global (CMIP5, CMIP6 and HighResMIP) and regional (CORDEX RCMs) model simulations at different resolutions (~300–12.5 km) and found that high-resolution models reduce the biases for extreme precipitation. They also found that the effect of horizontal resolutions for extreme precipitation is mostly in regions with complex topography and in the summer season when precipitation is mostly caused by convective processes, in agreement with Iles et al. (2020). The reduced biases in extreme precipitation near topography in high-resolution models is mostly due to an improved representation of topography, coastlines, and small-scale processes such as convection and diffusion. However, Strandberg & Lind (2021) showed that models with higher horizontal resolution overestimate the intensity of extreme precipitation in some regions over Europe. Moreover, α ... [1]

248 period 1979-2014 and are extended to 2019 using sea-surface temperature (SST) from ERA5  
 249 and the Shared Socioeconomic Pathway 5 (SSP5-8.5) scenario from CMIP6. OpenIFS  
 250 simulations use 91 vertical levels (L91) and the different horizontal resolutions: low resolution  
 251 (Tco95, ~100 km), medium resolution (Tco199, ~50 km), and high resolution (Tco399, ~25  
 252 km). For the low resolution, additional sensitivity experiments use different model time steps  
 253 i.e., 60, 30, and 15 minutes and we refer to these experiments as LR60m, LR30m, and LR,  
 254 respectively. For medium and high resolution, the same model time step is used (i.e., 15  
 255 minutes), of which experiments refer to as MR and HR, respectively. While the OpenIFS uses  
 256 a reduced octahedral grid (Malardel et al., 2016), the final output used in this study has been  
 257 interpolated to a regular grid using the XIOS output server. The LR, LR30m and LR60m data  
 258 were interpolated to a global 0.9° grid while the MR and HR data were interpolated to a global  
 259 0.45° grid, i.e., we are not investigating extreme precipitation in high resolution simulations in  
 260 their native grid, which will be investigated in future study. The simulations used here were  
 261 used by Savita et al. (2024) who found improvements in the surface zonal wind, Rossby wave  
 262 amplitude and phase speed, weather regime patterns, and surface-air temperature when  
 263 reducing a model time step from 60 minutes to 30 and 15 minutes in low resolution or  
 264 increasing the horizontal resolution from 100 km to 50 and 25 km. However, Savita et al. (2024)  
 265 did not find such improvement in the mean precipitation bias by increasing horizontal  
 266 resolution or reducing the model time step.

267 To validate OpenIFS simulations, we use the gridded daily precipitation observational data  
 268 from Global Precipitation Climatology Centre (GPCC) with resolution of  $1^\circ \times 1^\circ$  for the period  
 269 1982–2019 (Ziese et al., 2022) as well as the reanalysis data from the ECMWF Reanalysis v5  
 270 (ERA5) for 1979–2019 (Hersbach et al., 2023). ERA5 is based on the IFS Cy41r2, with 31 km  
 271 horizontal resolution and 137 levels (Hersbach et al., 2020). We analyzed total, large-scale,  
 272 and convective precipitation in this study. The total precipitation (convective plus large-scale  
 273 precipitation) in the IFS is the accumulated precipitation, comprising of rain and snow, that  
 274 falls to the Earth's surface, and it is not assimilated in the IFS. The convective precipitation is  
 275 generated by the convection scheme in the IFS, which represents convection at spatial scales  
 276 smaller than the grid box. The convection scheme follows Sundqvist (1978), which is also used  
 277 in the OpenIFS. The large-scale precipitation is generated by the cloud scheme (Khairoutdinov  
 278 & Kogan, 2000), which represents the formation and dissipation of clouds and large-scale  
 279 precipitation due to changes in atmospheric quantities (such as pressure, temperature, and  
 280 moisture) predicted directly by the IFS at spatial scales of the grid box or larger. The



281 autoconversion/accretion parameterization is a non-linear function of the mass of both liquid  
 282 cloud and rainwater. The calculation follows Khairoutdinov & Kogan (2000) which is derived  
 283 from large eddy simulation studies of drizzling stratocumulus clouds, and this scheme is also  
 284 used in OpenIFS. Several studies have evaluated the performance of ERA5 and found that the  
 285 total precipitation in ERA5 is performing well over the US (Tarek et al., 2020; Xu et al., 2019).  
 286 For global precipitation, the mean absolute difference over 50° S–50° N between ERA5 and  
 287 TRMM/3B43 is 0.58 mm/d; the global-mean correlation with GPCP data is 0.77, which is  
 288 better compared to ERA-Interim (0.63 mm/d and 0.67) (Hersbach et al., 2020). ERA5 also  
 289 performs well in polar regions in representing wind, temperature and humidity (Graham et al.,  
 290 2019; Tetzner et al., 2019; Wang et al., 2019).

291 Here we analyze daily ERA5 and the OpenIFS data over Europe (30° N–72° N, 10° W–40° E)  
 292 for the period of 1982–2019 to be consistent with GPCC dataset. For comparison, the ERA5,  
 293 GPCC, MR, and HR data are remapped to LR ( $\sim 0.9375^\circ \times 0.9375^\circ$ ) using the second-order  
 294 conservative remapping method. The second-order conservative method includes the gradient  
 295 across the source cell, which is not included in the first-order conservative method. Therefore,  
 296 it gives a smoother, more accurate representation of the source field (Jones, 1998).

## 297 298 **2.2 Methods**

### 299 **Calculation of $q^{\text{th}}$ percentile value**

300 We calculated different percentile values using total precipitation from GPCC, ERA5, and  
 301 OpenIFS simulations. When we calculated the  $q^{\text{th}}$  percentile value, the normalized ranking  
 302 usually did not match the location of the  $q^{\text{th}}$  percentile exactly, which means the  $q^{\text{th}}$  lies between  
 303 two indices. Therefore, we determined the location first, then computed the  $q^{\text{th}}$  value by  
 304 interpolating between the two nearest values based on the location. Here we used the formula  
 305 below to find the location:

$$306 \quad j = q \cdot (n-1) \quad (1)$$

307  $n$  is the length of the sample,  $q$  is the desired percentile,  $j$  is the location which is the distance  
 308 from the first value  $X_1$  ( $X_m$  are the sorted sample values,  $m=1, 2, \dots, n$ ). Then we took  $i$  as the  
 309 nearest (lower) integer of  $j$  to get the  $q^{\text{th}}$  value  $P(q)$  by interpolating.

$$310 \quad P(q) = X_i + (X_{i+1} - X_i) \cdot (j-i) \quad (2)$$

311 There are other methods to determine the location of  $q^{\text{th}}$  percentile (Hyndman & Fan, 1996),  
 312 but here we use the ‘linear’ one.

313

### The convective contribution to extreme precipitation

To calculate the contribution of convective precipitation to total precipitation for a percentile range, at each grid point we accumulated the convective precipitation on all days when the total precipitation is in that percentile range, then divided it by the accumulated total precipitation on those days to get the fraction of convective precipitation.

### Calculation of RMSE values

We used the root-mean-square error (RMSE) referenced to GPCC that measures the performance of ERA5 and OpenIFS simulations:

$$RMSE = \sqrt{\frac{\sum_{i=1}^n (x_{mi} - x_{oi})^2}{n}} \quad (3)$$

$x_{mi}$  is the value at  $i$  grid point for ERA5 or OpenIFS simulations,  $x_{oi}$  is the value for GPCC,  $n$  is the number of land grid points over Europe. Using equation (3), we calculated the RMSE values for different percentile ranges. Smaller RMSE values mean the biases between OpenIFS (or ERA5) and GPCC are smaller i.e., the model simulations and ERA5 are performing better.

### Confidence intervals

We calculated the 2.5 to 97.5<sup>th</sup> confidence intervals (CI) for the RMSE for each percentile with a bootstrap method. For example, to calculate the CI for the RMSE of HR (referenced to GPCC observation), we randomly chose  $n$  grid cell pairs from GPCC and HR over European land, then calculated their RMSE ( $n$  is the number of total land grid points over Europe). This process was repeated for 2000 times. We took the 2.5<sup>th</sup> and 97.5<sup>th</sup> percentiles of the distribution of the 2000 RMSEs as the 95 % CI. If the CI for different simulations do not overlap then we refer that they are significantly different.

## 3. Results

### 3.1 Extreme precipitation over Europe

We show the time series of 99<sup>th</sup> percentile precipitation calculated from all grid points and all days in each year over the period 1982–2019 from GPCC, ERA5, and OpenIFS simulations over Europe (Fig. 1). The ERA5 simulates an inter-annual variability of the 99<sup>th</sup> percentile precipitation similar to that in GPCC. For example, the peak in 2010 and the low in 1994 are well reproduced in the ERA5. OpenIFS simulations do not reproduce the same inter-annual

variability as in GPCC or ERA5 but LR and HR do reproduce the 95 % significant positive trend observed in GPCC (0.03 mm/d/y, not shown), which are  $\sim 0.2$  mm/d/y for both LR and HR, and it is not significant for MR. We note that the OpenIFS simulations use observed SST and sea-ice concentrations as boundary conditions, but ozone is taken from a photochemical equilibrium (Cariolle & Teyss  re, 2007) and aerosol concentrations are taken from Copernicus Atmosphere Monitoring Service (CAMS) monthly climatology. Therefore, we do not expect LR, MR and HR to reproduce trends driven by ozone or aerosols forcing. We also find that both ERA5 and OpenIFS simulations have relatively lower 99<sup>th</sup> percentile precipitation rates compared to GPCC (Fig. 1). The RMSE for ERA5 (0.36 mm/d) is lower than for OpenIFS simulations which is largest for LR (2.03 mm/d) and decreases with increasing horizontal resolution (i.e., 1.13 mm/d for MR and 0.69 mm/d for HR). Note that Fig. 1 does not contain any spatial information and that a mismatch between model data and observations can be due to the 99<sup>th</sup> percentile occurring in different regions and/or with different magnitudes. The RMSE analysis suggests that ERA5 and HR are close to GPCC and LR is far from GPCC.

Figure 2a–e shows the spatial distribution of the 99<sup>th</sup> percentile precipitation over Europe for all days in each season for all years in GPCC, ERA5, and OpenIFS simulations, respectively. In general, the extreme precipitation is very low ( $\sim 2$  mm/d) in Northern Africa, which is to be expected since the mean precipitation is only 0.5 mm/d in those regions (Fig. S1). The extreme precipitation exceeds 30 mm/d over mountain areas (e.g., Scandinavian mountains, Alps, and Iberian Peninsula) and the north coast of the Mediterranean but is otherwise lower ( $\sim 15$  mm/d). The spatial distribution of extreme precipitation matches that of the mean precipitation pattern (Fig. S1). The high 99<sup>th</sup> percentile precipitation near mountains is likely due to the forced ascent of westerly (Scandinavia, Iberian Peninsula, British Isles) and southerly (Alps) winds. The high 99<sup>th</sup> precipitation in the north of the Mediterranean is likely because of warm and moist southerly winds from the Mediterranean Sea. The ERA5 and OpenIFS simulations overall reproduce the spatial distribution of the 99<sup>th</sup> percentile precipitation from GPCC. However, the magnitudes are different, particularly over the Scandinavian mountains, the Alps, and central Europe near 50° N (Fig. 2a–e). Figure 2f–i show the regional biases for the 99<sup>th</sup> percentile precipitation referenced to GPCC. LR mostly underestimates the 99<sup>th</sup> percentile precipitation in mountainous areas and deserts by more than 25 % (Fig. 2g) and the biases are reduced when horizontal resolution is increased in MR and HR (Fig. 2h–i). We also notice that LR

underestimates the 99<sup>th</sup> percentile precipitation south of the Alps but overestimates it to the north (Fig. 2 (g)), whereas this bias is negligible in higher-resolution simulations (Fig. 2h-i). This could be because the moist southerly winds do not ascend high enough with LR, therefore there is less precipitation formed on the southern side and more moisture is advected over the mountain. The reduced biases near mountain regions in the higher-resolution simulations are likely because higher resolution has a better representation of topography and vertical velocity. A cross section of the topography and annual-mean vertical velocity at 850 hPa and 62° N (Fig. S2 and S3) highlight that the higher-resolution simulations resolve steeper topography, which leads to more ascent and thus more precipitation.

The 99<sup>th</sup> percentile precipitation over the Alps is more realistic with higher horizontal resolution compared to lower resolution. However, all simulations as well as ERA5 exhibit a negative bias over northeast Italy and west Slovenia (Fig. 2f-i). By analyzing the EOBS data, we find a similar negative bias as in GPCC, but a positive bias in GPCP (Fig. S8d-k). We notice that the extreme precipitation over Alps (include Slovenia) in GPCP is lower than GPCC and EOBS (Fig. S8a-c), which is likely due to the different data sources and grid methods in different observation datasets (e.g., GPCC and EOBS are gauge-based gridded data on the land, but GPCP data combines microwave and infrared measurements, satellites and rain gauges). We do not know which observation dataset is more realistic, therefore, the cause of the negative bias near Slovenia could be a bias in GPCC or a persistent model bias in the ECMWF-IFS on which both ERA5 and OpenIFS are based. In general, ERA5 has a lower RMSE (2.6 mm/d) for extreme total precipitation than OpenIFS simulations, i.e., ERA5 has overall lower biases than LR (4.0 mm/d) and is similar to MR (3.0 mm/d) and HR (2.9 mm/d).

We next calculate the trend for the annual 99<sup>th</sup> percentile precipitation over Europe (Fig. S4 & S5) and find that the 99<sup>th</sup> percentile precipitation has a large positive trend in central Europe and a negative trend to the north of the Alps in GPCC (Fig. S4a). The ERA5 reproduces the pattern of the trend found in GPCC but not significant. However, OpenIFS simulations do not have consistent patterns with GPCC (Fig. S4c-e, Fig. S5c-e), with only LR60m reproducing the large positive trend in central Europe (Fig. S5c). Overall, the trend is largely underestimated over central Europe but overestimated over northern Europe in OpenIFS simulations. We have not found any consistent improvement across the horizontal resolution and model time step.

**Deleted:** Lavers et al. (2022) also found too much extreme precipitation on the north side of the Alps in ERA5 during a storm. ...

**Deleted:** T

In addition to the 99<sup>th</sup> percentile precipitation, we calculate annual total precipitation in different percentile ranges, such as 70<sup>th</sup>–80<sup>th</sup>, 80<sup>th</sup>–90<sup>th</sup>, 90<sup>th</sup>–95<sup>th</sup>, 95<sup>th</sup>–99<sup>th</sup>, 99<sup>th</sup>–99.5<sup>th</sup>, 99.5<sup>th</sup>–99.9<sup>th</sup> and larger than 99.9<sup>th</sup> (i.e., >99.9<sup>th</sup>) percentile. We calculate the RMSEs for ERA5 and OpenIFS simulations referenced to GPCC in each range and find that the RMSEs for ERA5 and OpenIFS simulations vary strongly across percentile ranges (Fig. 3). The RMSEs increase exponentially with increasing percentiles, from less than 1 mm/d at the 70<sup>th</sup>–80<sup>th</sup> percentile range to ~8 mm/d above the 99.9<sup>th</sup> percentile range. The largest RMSE is found for LR60m above the 99.9<sup>th</sup> percentile range which is around 12 mm/d [CI: 11.3–12.8 mm/d]. We also find that the RMSEs decrease with finer horizontal resolution for all percentile ranges. The CI of the RMSEs from LR do not overlap with those from higher horizontal resolutions for any percentile range, i.e., the biases from LR are significantly different from that at higher resolutions and thus clearly sensitive to the horizontal resolution. We also find that the RMSE differences between LR simulation and the higher-resolution simulations as well as ERA5 are larger at higher percentile ranges (>95<sup>th</sup>) than those at lower percentile ranges (<95<sup>th</sup>). Thus, we conclude that extreme precipitation is more sensitive to horizontal resolution than precipitation at lower percentile ranges (<95<sup>th</sup>). ERA5 has the smallest RMSE of all datasets above the 95<sup>th</sup> percentile ranges, i.e., ERA5 has a better representation of the extreme precipitation than our OpenIFS simulations (Fig 3).

The RMSEs for LR60m, LR30m, and LR are increasing with increasing model time steps. However, the CI of RMSE overlap at all percentile ranges, i.e., the sensitivity of precipitation to the model time step is not statistically significant in the low-resolution configurations. While the model time step may influence precipitation, especially convective precipitation, errors from poorly resolved topography probably have a large impact on the RMSE, which would explain the lack of sensitivity to the model time step.

### 3.2 Relative roles of convective and large-scale precipitation

Total precipitation is the sum of convective and large-scale precipitation. Convective precipitation is related with not explicitly resolved convective motions. Their scales are smaller than the resolution of the model, therefore need to be parametrized. With increasing resolution, convective precipitation decreases as higher-resolution simulations can resolve smaller scales of vertical motions. On the other hand, large-scale precipitation is likely to originate from large-scale synoptic storms. As horizontal resolution increases, grid cells become smaller and

452 humidity saturation level is reached faster, thus, large-scale precipitation increases (Hertwig et  
453 al., 2015). As we showed in Fig. S7, the large-scale precipitation is more frequent in MR and  
454 HR compared to LR when it is higher than 20 mm/d. Higher resolution simulations can produce  
455 higher large-scale precipitation better than LR. However, convective precipitation also  
456 increases with increasing resolution for convective precipitation higher than 20 mm/d (Fig. S6),  
457 but if we only consider the European area-mean values (not shown), the JJA convective  
458 precipitation decreases and the large-scale precipitation increases with increasing resolution,  
459 which is consistent with Hertwig et al. (2015) result. We find that the convective precipitation  
460 is sensitive to time step and the smallest time step has the minimum convective precipitation.  
461 The large-scale precipitation is insensitive to time step during JJA and DJF.

462  
463 We calculate the fractions of convective and large-scale precipitation in total precipitation for  
464 days when the total precipitation exceeds the 99<sup>th</sup> percentile in all model simulations and ERA5  
465 (Fig. 4 & 5). Note that, GPCC does not provide convective and large-scale precipitation  
466 separately, therefore we compare our OpenIFS simulations to the ERA5 dataset to assess the  
467 effect of resolution and model time step in the model simulations. We note that ERA5 is a  
468 reanalysis dataset where precipitation is a parametrized variable, and observations of which are  
469 not assimilated over Europe. ERA5 monthly precipitation has a good agreement with GPCC  
470 on the land, with correlations above 90 % for most of Europe, and above 70 % for Australia,  
471 Asia, and North America (Bell et al., 2021). ERA5 also shows smaller biases for mean  
472 precipitation than other reanalysis datasets in the tropics compared to the Global Precipitation  
473 Climatology Project (GPCP), with relative biases of 13 % for ERA5, 17 % for MERRA-2 and  
474 36 % for JRA-55 (Hassler & Lauer, 2021). The biases for mean precipitation are found smaller  
475 over extra-tropics than the tropics compared to the gauge-based precipitation observations,  
476 particularly agreeing well with observations over central Europe and South Asia (Lavers et al.,  
477 2022). Moreover, ERA5 can capture the locations and patterns of highest precipitations in  
478 observations, but cannot simulate the magnitude (Lavers et al., 2022). We also find that the  
479 extreme precipitation over Europe in ERA5 is closer to observations than models (Fig. 1, 2,  
480 and 3), therefore, we use ERA5 for the benching mark here although it has some known biases.

481  
482 The ERA5 data and OpenIFS simulations show that, in DJF, the extreme precipitation is nearly  
483 100 % large-scale precipitation over northern Europe, more than 90 % over central Europe,  
484 and more than 70 % over western and southern Europe (Fig. 5a–d). However, in JJA, large-  
485 scale precipitation makes up most of the extreme precipitation over northern Europe (>70 %)

Deleted: realism

Deleted: of

488 while convective precipitation makes up most of the extreme precipitation in the Mediterranean  
 489 region (>70 %) (Fig. 4a–d). The OpenIFS simulations largely reproduce the pattern of the  
 490 fraction of convective precipitation found in ERA5, but we note differences in magnitudes (Fig.  
 491 4e–g, and Fig. 5e–g)). In JJA, the OpenIFS simulates the contribution of the convective  
 492 precipitation ~~very close to ERA5~~ over Scandinavia where the extreme precipitation is mostly  
 493 large-scale precipitation, but ~~has more convective precipitation~~ for other areas over Europe  
 494 ~~compared to ERA5~~ (Fig. 4e–g). The RMSEs from MR (0.10 mm/d [CI: 0.09–0.10 mm/d]) and  
 495 HR (~0.09 mm/d [CI: 0.09–0.10 mm/d]) are not significantly different, while LR (~0.12 mm/d  
 496 [CI: 0.12–0.13 mm/d]) is significantly larger than those in MR and HR. In DJF, the OpenIFS  
 497 ~~has less~~ contribution from convective precipitation ~~than ERA5~~ except for near-coastal areas  
 498 (Fig. 5e–f). That is, the contribution from large-scale precipitation is ~~more in OpenIFS than~~  
 499 ~~ERA5~~, and their ~~difference~~ is reduced with higher horizontal resolution, i.e., in MR and HR.

500  
 501 Further, we explore the relative roles of horizontal resolution and time step for the large-scale  
 502 and convective precipitation at different percentile ranges (Fig. 6). In general, the RMSEs  
 503 increase with increasing percentiles, but also decrease with increasing horizontal resolution and  
 504 shorter model time step. The RMSE reduces for higher percentile in higher resolution due to  
 505 better representation of topography, and in smaller model time step due to enhanced convection.  
 506 The exceptions are the total precipitation above the 99.5<sup>th</sup> percentile in JJA where the RMSEs  
 507 from LR are larger than LR30m (Fig. 6a), and the convective precipitation above the 99<sup>th</sup>  
 508 percentile in JJA and DJF where the RMSEs from HR are larger than MR (Fig. 6c & f).

509  
 510 The CI for RMSEs of total precipitation from LR, MR and HR in DJF and JJA do not overlap  
 511 for all percentile ranges, thus there is a significant sensitivity of the total precipitation to the  
 512 horizontal resolution, particularly for extreme total precipitation. The exceptions are the total  
 513 precipitation below the 90<sup>th</sup> percentile ranges and above the 99.9<sup>th</sup> percentile range in JJA  
 514 where the CI for RMSEs in MR and HR overlap (Fig. 6a). However, the sensitivity is not found  
 515 for the global-mean total precipitation by increasing horizontal resolution (Savita et al., 2024).  
 516 For the large-scale precipitation in JJA, the CI for RMSEs from LR do not overlap with those  
 517 from MR and HR at higher percentile ranges (>95<sup>th</sup>), but overlap at lower percentile ranges  
 518 (<95<sup>th</sup>) (Fig. 6b). That is, the large-scale precipitation from the extreme precipitation is  
 519 sensitive to the horizontal resolution. We note that a reduced ~~RMSE with increasing resolutions~~  
 520 ~~at >99<sup>th</sup>~~ is not found for the convective precipitation in JJA (Fig. 6c), and we conclude that the

Deleted: quite well

Deleted: overestimates

Deleted: that

Deleted: underestimates the

Deleted: overestimated

Deleted: bias

Deleted: bias

Formatted: Superscript

horizontal resolution dependence of extreme total precipitation in JJA comes from the large-scale precipitation. For DJF, the large-scale precipitation is sensitive to the horizontal resolution for all percentile ranges, where the CI for RMSEs in LR do not overlap with those from MR and HR (Fig. 6e). The convective precipitation in DJF is also sensitive to the horizontal resolution (Fig. 6f), however there is little convection precipitation in DJF, thus the sensitivity for convective precipitation in DJF is not important. Therefore, the resolution dependence of extreme total precipitation is mostly dominated by the large-scale precipitation in DJF.

For the model time step, the CI for RMSEs of total precipitation from LR60m, LR30m, and LR overlap at all percentile ranges in both JJA and DJF (Fig. 6a & d), i.e., the extreme total precipitation is not sensitive to the model time step in a significant way in the low-resolution simulations. Similarly, the mean total precipitation is also found insensitive to the model time step (Savita et al., 2024). Both the large-scale and convective precipitation are sensitive to the model time step particularly above the 95<sup>th</sup> percentile ranges in JJA (Fig. 6b & c). The convective precipitation is more sensitive to the model time step than the large-scale precipitation in JJA, but in DJF the sensitivity is found only for the large-scale precipitation (Fig. 6e). Also, the lack of sensitivity for convective precipitation in DJF may be because there is almost no convective precipitation in DJF.

#### 4. Discussion and Conclusion

We have investigated the sensitivity of extreme precipitation across different horizontal resolutions and model time steps in atmosphere-only experiments with the OpenIFS. Comparing extreme precipitation (defined as total daily precipitation at the 99<sup>th</sup> percentile) from OpenIFS simulations, reanalysis (ERA5), and observation (GPCC), we find that MR and HR mostly better represent the precipitation extremes compared to LR. We also found a more significant sensitivity to the horizontal resolution for the precipitation above the 95<sup>th</sup> percentile and less sensitivity for lower percentile ranges (<95<sup>th</sup>) (Fig. 3). These OpenIFS-based results are similar to Kopparla et al. (2013), who found that the bias of extreme precipitation in the high-resolution simulation (25 km) is reduced compared to the lower-resolution simulations (100 km and 200 km) over Europe in their atmospheric model, but not for precipitation at lower percentiles (i.e., <95<sup>th</sup>). However, the sensitivity to the horizontal resolution found by Kopparla et al. (2013) was not significant over Europe which is rather different from our results as we have found a significant difference across the horizontal resolutions. In contrast to the extreme



precipitation, the bias for global mean precipitation is not decreasing when increasing horizontal resolution from ~200 km to ~100 km or ~50 km in the ECHAM6-AMIP simulations (Hertwig et al., 2015), and also in other GCMs (e.g., OpenIFS, HadGEM1 and HadGEM3) (Demory et al., 2020; Savita et al., 2024; Schiemann et al., 2014). However, Delworth et al. (2012) found an improvement in the global mean precipitation with increasing horizontal resolution in a coupled model (GFDL).

The improvements due to increasing horizontal resolution for the extreme precipitation are mostly over the mountain areas, consistent with previous studies which found the effect of horizontal resolution being largest in areas with complex topography over Europe and also other regions for mean and extreme precipitation (Demory et al., 2020; Iles et al., 2020; Monerie et al., 2020; Prein et al., 2013; Torma et al., 2015). The sensitivity to the horizontal resolution comes from the large-scale precipitation, which is likely because of the better-resolved topography. However, the convective precipitation in JJA is more sensitive to the model time step than it is to the horizontal resolution, likely because there is an increase in shallow and mid-level convection with a shorter time step in the OpenIFS (Savita et al., 2024), thus we get more convective precipitation.

In our results, larger improvements are obtained when the horizontal resolution is increased from LR to MR, but relatively smaller improvements from MR to HR. Similar results are also found in Roberts et al. (2018), where the climatological surface biases are relatively insensitive when increasing the atmospheric resolution from ~50 km to ~25 km in the ECMWF-IFS. Jung et al. (2012) also showed that the largest improvements in extratropical cyclones, Euro-Atlantic blocking, tropical mean precipitation, and tropospheric circulation are found when increasing horizontal resolution from 126 km to 39 km with relatively small further changes from 39 km to 16 and to 10 km in ECMWF atmosphere model. Kopparla et al. (2013) and Bacmeister et al. (2014) found much improvement for the mean precipitation and extreme precipitation with the increasing atmospheric resolution from ~200 km to 100 km, but less improvement from ~100 km to ~25 km. It is likely due to a lack of tuning with the changing horizontal resolution. The above conclusions are valid over Europe, but maybe not valid for other regions such as the tropics and subtropics, where the extreme precipitation is often triggered by tropical cyclones (Gori et al., 2022; Zhu & Quiring, 2022). The distribution of tropical cyclones intensity and the structure of the most intense storm can be adequately simulated at 16 and 10-km horizontal resolutions, but not at 126 and 39-km resolutions in ECMWF-IFS (Manganello et al., 2011).

596  
597  
598 Moreover, the choice of observation dataset is a key factor for assessing the impact of the  
599 horizontal resolution and model time step on extreme precipitation. Most observation  
600 precipitation data are from one of the three categories, gauge-based products, satellite products,  
601 and merged satellite-gauge products. Since the satellite products are constructed with satellite  
602 microwave and/ or infrared measurements, with/ without gauged-adjusted estimates,  
603 differences exist between these products. Besides, the gauge-based products are highly  
604 dependent on the choice of stations and interpolation schemes. It is hard to say which product  
605 is closer to reality, as different regions may have different observation datasets that suit best  
606 for the analysis. In particular, we note that not all products are suitable for extreme analysis.  
607 For example, GPCP's main scope is to construct a reliable climate data record and has been  
608 developed with a priority of ensuring the long-term stability of data (Adler et al., 2017).  
609 Masunaga et al. (2019) found that the frequency of GPCP daily precipitation quickly drops  
610 below all other datasets once the precipitation exceeds 30 mm/d. Also, the time series of GPCP  
611 extreme precipitation over the ocean exhibits a jump to lower 99<sup>th</sup> percentiles in late 2008/early  
612 2009 which is not present in all other datasets, coinciding with the change in utilization of  
613 SSM/I and SSMIS. The lower 99<sup>th</sup> precipitation suggests that the GPCP dataset might not be  
614 applied to extreme analysis (Masunaga et al., 2019). Therefore, we only use GPCC observation  
615 data as the reference to explore the model performance. In Fig. 2f–i the 99<sup>th</sup> percentile  
616 precipitation is largely underestimated in the eastern Alp region by ERA5 and all model  
617 simulations. The biases are insensitive to horizontal resolution. It is likely a persistent model  
618 bias in the ECMWF-IFS or a bias in GPCC. Analyzing multiple precipitation products instead  
619 of relying on a single one may be a good way to reduce these biases.

620

## 621 **Code and data variability**

622 The OpenIFS model requires a software license agreement with ECMWF to use it, and  
623 OpenIFS's license is easily given free of charge to any academic or research institute. The  
624 details of OpenIFS are available at  
625 <https://confluence.ecmwf.int/display/OIFS/About+OpenIFS> (ECMWF, 2018). We used the  
626 same simulation that used in Savita et al. (2024) and therefore do not provide the data needed  
627 to reproduce the simulations here. All data (runscripts, input data etc) needed to reproduce the  
628 simulations can be found in Savita et al. (2024) in code and data variability section. The jupyter  
629 notebook scripts used in this study to produce the plots can be found at

**Deleted:** The above conclusions are valid over Europe, but also valid for other regions such as the tropics and subtropics. For example, the predictions of tropical cyclone intensity are markedly improved when the horizontal resolution of the atmosphere model is increased from 120 km to 40 km, but not for 15 to 10 km (Jung et al., 2012), which often triggers extreme precipitation (Gori et al., 2022; Zhu & Quiring, 2022)....

638 <https://doi.org/10.5281/zenodo.10887652>. The raw model output is available from the authors  
639 upon reasonable request. The observation and reanalysis datasets used in this study can be  
640 downloaded from GPCC ([https://opendata.dwd.de/climate\\_environment/GPCC/html/fulldata-](https://opendata.dwd.de/climate_environment/GPCC/html/fulldata-daily_v2022_doi_download.html)  
641 [daily\\_v2022\\_doi\\_download.html](https://opendata.dwd.de/climate_environment/GPCC/html/fulldata-daily_v2022_doi_download.html), Ziese et al., 2022) and ERA5  
642 (<https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels?tab=form>,  
643 Hersbach et al., 2023).

644  
645 **Authors contributions.** AS and JK conducted all the OpenIFS simulations. YL did the  
646 analysis and writing with substantial contribution from JK, AS and WP.

647  
648 **Competing interests.** The contact author has declared that none of the authors has any  
649 competing interests.

650  
651 **Acknowledgements.** Yingxue Liu is supported by China Scholarship Council (CSC, grant no.  
652 202004910401). Joakim Kjellsson and Abhishek Savita are supported by JPI Climate/Ocean  
653 (ROADMAP project, grant no. 01LP2002C). Wonsun Park was supported by IBS (grant no.  
654 IBS-R028-D1). We thank the OpenIFS team at ECMWF for the technical support. All the  
655 OpenIFS simulations were conducted on the HLRN machine under shk00018 project resources.  
656 All the analysis and data storage were conducted on computer clusters at GEOMAR and Kiel  
657 University Computing Center (NESH).

658  
659 **Financial support.** This research is financially supported by CSC (grant no. 202004910401)  
660 and ROADMAP project (grant no. 01LP2002C).

661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671

672

## 673 References

- 674 Adler, R. F., Gu, G., Sapiiano, M., Wang, J. J., & Huffman, G. J. (2017). Global  
675 Precipitation: Means, Variations and Trends During the Satellite Era (1979–2014).  
676 In *Surveys in Geophysics* (Vol. 38, Issue 4, pp. 679–699). Springer Netherlands.  
677 <https://doi.org/10.1007/s10712-017-9416-4>
- 678 Alexander, L. V., Fowler, H. J., Bador, M., Behrangi, A., Donat, M. G., Dunn, R., Funk,  
679 C., Goldie, J., Lewis, E., Rogé, M., Seneviratne, S. I., & Venugopal, V. (2019). On  
680 the use of indices to study extreme precipitation on sub-daily and daily timescales.  
681 *Environmental Research Letters*, 14(12). <https://doi.org/10.1088/1748-9326/ab51b6>
- 682 Avila, F. B., Dong, S., Menang, K. P., Rajczak, J., Renom, M., Donat, M. G., &  
683 Alexander, L. V. (2015). Systematic investigation of gridding-related scaling  
684 effects on annual statistics of daily temperature and precipitation maxima: A case  
685 study for south-east Australia. *Weather and Climate Extremes*, 9, 6–16.  
686 <https://doi.org/10.1016/j.wace.2015.06.003>
- 687 Bacmeister, J. T., Wehner, M. F., Neale, R. B., Gettelman, A., Hannay, C., Lauritzen, P.  
688 H., Caron, J. M., & Truesdale, J. E. (2014). Exploratory high-resolution climate  
689 simulations using the community atmosphere model (CAM). *Journal of Climate*,  
690 27(9), 3073–3099. <https://doi.org/10.1175/JCLI-D-13-00387.1>
- 691 Bador, M., Boé, J., Terray, L., Alexander, L. V., Baker, A., Bellucci, A., Haarsma, R.,  
692 Koenigk, T., Moine, M. P., Lohmann, K., Putrasahan, D. A., Roberts, C., Roberts,  
693 M., Scoccimarro, E., Schiemann, R., Seddon, J., Senan, R., Valcke, S., & Vanniere,  
694 B. (2020). Impact of Higher Spatial Atmospheric Resolution on Precipitation  
695 Extremes Over Land in Global Climate Models. *Journal of Geophysical Research:*  
696 *Atmospheres*, 125(13). <https://doi.org/10.1029/2019JD032184>
- 697 Barrett, A. I., Wellmann, C., Seifert, A., Hoose, C., Vogel, B., & Kunz, M. (2019). One  
698 Step at a Time: How Model Time Step Significantly Affects Convection-Permitting  
699 Simulations. *Journal of Advances in Modeling Earth Systems*, 11(3), 641–658.  
700 <https://doi.org/10.1029/2018MS001418>
- 701 Bell, B., Hersbach, H., Simmons, A., Berrisford, P., Dahlgren, P., Horányi, A., Muñoz-  
702 Sabater, J., Nicolas, J., Radu, R., Schepers, D., Soci, C., Villaume, S., Bidlot, J. R.,  
703 Haimberger, L., Woollen, J., Buontempo, C., & Thépaut, J. N. (2021). The ERA5  
704 global reanalysis: Preliminary extension to 1950. *Quarterly Journal of the Royal*  
705 *Meteorological Society*, 147(741), 4186–4227. <https://doi.org/10.1002/qj.4174>
- 706 Caldwell, P. (2010). California wintertime precipitation bias in regional and global  
707 climate models. *Journal of Applied Meteorology and Climatology*, 49(10), 2147–  
708 2158. <https://doi.org/10.1175/2010JAMC2388.1>
- 709 Cariolle, D., & Teyssède, H. (2007). Atmospheric Chemistry and Physics A revised  
710 linear ozone photochemistry parameterization for use in transport and general  
711 circulation models: multi-annual simulations. In *Atmos. Chem. Phys* (Vol. 7).  
712 [www.atmos-chem-phys.net/7/2183/2007/](http://www.atmos-chem-phys.net/7/2183/2007/)
- 713 Chan, S. C., Kendon, E. J., Fowler, H. J., Blenkinsop, S., Ferro, C. A. T., & Stephenson,  
714 D. B. (2013). Does increasing the spatial resolution of a regional climate model  
715 improve the simulated daily precipitation? *Climate Dynamics*, 41(5–6), 1475–1495.  
716 <https://doi.org/10.1007/s00382-012-1568-9>
- 717 Delworth, T. L., Rosati, A., Anderson, W., Adcroft, A. J., Balaji, V., Benson, R., Dixon,  
718 K., Griffies, S. M., Lee, H. C., Pacanowski, R. C., Vecchi, G. A., Wittenberg, A.  
719 T., Zeng, F., & Zhang, R. (2012). Simulated climate and climate change in the


- GFDL CM2.5 high-resolution coupled climate model. *Journal of Climate*, 25(8), 2755–2781. <https://doi.org/10.1175/JCLI-D-11-00316.1>
- Demory, M. E., Berthou, S., Fernández, J., Sørland, S. L., Brogli, R., Roberts, M. J., Beyerle, U., Seddon, J., Haarsma, R., Schär, C., Buonomo, E., Christensen, O. B., Ciarlo, J. M., Fealy, R., Nikulin, G., Peano, D., Putrasahan, D., Roberts, C. D., Senan, R., ... Vautard, R. (2020). European daily precipitation according to EURO-CORDEX regional climate models (RCMs) and high-resolution global climate models (GCMs) from the High-Resolution Model Intercomparison Project (HighResMIP). *Geoscientific Model Development*, 13(11), 5485–5506. <https://doi.org/10.5194/gmd-13-5485-2020>
- Gori, A., Lin, N., Xi, D., & Emanuel, K. (2022). Tropical cyclone climatology change greatly exacerbates US extreme rainfall–surge hazard. *Nature Climate Change*, 12(2), 171–178. <https://doi.org/10.1038/s41558-021-01272-7>
- Graham, R. M., Hudson, S. R., & Maturilli, M. (2019). Improved Performance of ERA5 in Arctic Gateway Relative to Four Global Atmospheric Reanalyses. *Geophysical Research Letters*, 46(11), 6138–6147. <https://doi.org/10.1029/2019GL082781>
- Hassler, B., & Lauer, A. (2021). Comparison of reanalysis and observational precipitation datasets including era5 and wfde5. *Atmosphere*, 12(11). <https://doi.org/10.3390/atmos12111462>
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., ... Thépaut, J. N. (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999–2049. <https://doi.org/10.1002/qj.3803>
- Hertwig, E., von Storch, J. S., Handorf, D., Dethloff, K., Fast, I., & Krismer, T. (2015). Effect of horizontal resolution on ECHAM6-AMIP performance. *Climate Dynamics*, 45(1–2), 185–211. <https://doi.org/10.1007/s00382-014-2396-x>
- Hyndman, R. J., & Fan, Y. (1996). Sample Quantiles in Statistical Packages. In *Source: The American Statistician* (Vol. 50, Issue 4).
- Iles, C. E., Vautard, R., Strachan, J., Joussaume, S., Eggen, B. R., & Hewitt, C. D. (2020). The benefits of increasing resolution in global and regional climate simulations for European climate extremes. *Geoscientific Model Development*, 13(11), 5583–5607. <https://doi.org/10.5194/gmd-13-5583-2020>
- Intergovernmental Panel on Climate Change. (2023). Weather and Climate Extreme Events in a Changing Climate. In *Climate Change 2021 – The Physical Science Basis* (pp. 1513–1766). Cambridge University Press. <https://doi.org/10.1017/9781009157896.013>
- Jones, P. W. (1998). *First-and Second-Order Conservative Remapping Schemes for Grids in Spherical Coordinates*.
- Jong, B. T., Delworth, T. L., Cooke, W. F., Tseng, K. C., & Murakami, H. (2023). Increases in extreme precipitation over the Northeast United States using high-resolution climate model simulations. *Npj Climate and Atmospheric Science*, 6(1). <https://doi.org/10.1038/s41612-023-00347-w>
- Jung, T., Miller, M. J., Palmer, T. N., Towers, P., Wedi, N., Achuthavarier, D., Adams, J. M., Altshuler, E. L., Cash, B. A., Kinter, J. L., Marx, L., Stan, C., & Hodges, K. I. (2012). High-resolution global climate simulations with the ECMWF model in project athena: Experimental design, model climate, and seasonal forecast skill. *Journal of Climate*, 25(9), 3155–3172. <https://doi.org/10.1175/JCLI-D-11-00265.1>
- Khairoutdinov, M., & Kogan, Y. (2000). *A New Cloud Physics Parameterization in a Large-Eddy Simulation Model of Marine Stratocumulus*.

- Kopparla, P., Fischer, E. M., Hannay, C., & Knutti, R. (2013). Improved simulation of extreme precipitation in a high-resolution atmosphere model. *Geophysical Research Letters*, 40(21), 5803–5808. <https://doi.org/10.1002/2013GL057866>
- Laprise, R. (2008). Regional climate modelling. *Journal of Computational Physics*, 227(7), 3641–3666. <https://doi.org/10.1016/j.jcp.2006.10.024>
- Lavers, D. A., Simmons, A., Vamborg, F., & Rodwell, M. J. (2022). An evaluation of ERA5 precipitation for climate monitoring. *Quarterly Journal of the Royal Meteorological Society*, 148(748), 3152–3165. <https://doi.org/10.1002/qj.4351>
- Li, C., Zwiers, F., Zhang, X., Li, G., Sun, Y., & Wehner, M. (2021). Changes in Annual Extremes of Daily Temperature and Precipitation in CMIP6 Models. *Journal of Climate*, 34, 3441–3460. <https://doi.org/10.1175/JCLI-D-19>
- Malardel, S., Wedi, N., Deconinck, W., & Kühnlein, C. (2016). *A new grid for the IFS*. <https://www.researchgate.net/publication/297695132>
- [Manganello, J. V., Hodges, K. I., Kinter, J. L., Cash, B. A., Marx, L., Jung, T., Achuthavarier, D., Adams, J. M., Altshuler, E. L., Huang, B., Jin, E. K., Stan, C., Towers, P., & Wedi, N. \(2012\). Tropical cyclone climatology in a 10-km global atmospheric GCM: Toward weather-resolving climate modeling. \*Journal of Climate\*, 25\(11\), 3867–3893. <https://doi.org/10.1175/JCLI-D-11-00346.1>](#)
- Masunaga, H., Schröder, M., Furuzawa, F. A., Kummerow, C., Rustemeier, E., & Schneider, U. (2019). Inter-product biases in global precipitation extremes. *Environmental Research Letters*, 14(12). <https://doi.org/10.1088/1748-9326/ab5da9>
- Mishra, S. K., & Sahany, S. (2011). Effects of time step size on the simulation of tropical climate in NCAR-CAM3. *Climate Dynamics*, 37(3), 689–704. <https://doi.org/10.1007/s00382-011-0994-4>
- Monerie, P. A., Chevuturi, A., Cook, P., Klingaman, N. P., & Holloway, C. E. (2020). Role of atmospheric horizontal resolution in simulating tropical and subtropical South American precipitation in HadGEM3-GC31. *Geoscientific Model Development*, 13(10), 4749–4771. <https://doi.org/10.5194/gmd-13-4749-2020>
- Myhre, G., Alterskjær, K., Stjern, C. W., Hodnebrog, M., Marelle, L., Samset, B. H., Sillmann, J., Schaller, N., Fischer, E., Schulz, M., & Stohl, A. (2019). Frequency of extreme precipitation increases extensively with event rareness under global warming. *Scientific Reports*, 9(1). <https://doi.org/10.1038/s41598-019-52277-4>
- O’Gorman, P. A. (2015). Precipitation Extremes Under Climate Change. In *Current Climate Change Reports* (Vol. 1, Issue 2, pp. 49–59). Springer. <https://doi.org/10.1007/s40641-015-0009-3>
- Prein, A. F., Gobiet, A., Suklitsch, M., Truhetz, H., Awan, N. K., Keuler, K., & Georgievski, G. (2013). Added value of convection permitting seasonal simulations. *Climate Dynamics*, 41(9–10), 2655–2677. <https://doi.org/10.1007/s00382-013-1744-6>
- Rauscher, S. A., O’Brien, T. A., Piani, C., Coppola, E., Giorgi, F., Collins, W. D., & Lawston, P. M. (2016). A multimodel intercomparison of resolution effects on precipitation: simulations and theory. *Climate Dynamics*, 47(7–8), 2205–2218. <https://doi.org/10.1007/s00382-015-2959-5>
- Savita, A., Kjellsson, J., Kedzierski, R. P., Latif, M., Rahm, T., Wahl, S., & Park, W. (2024). Assessment of climate biases in OpenIFS version 43r3 across model horizontal resolutions and time steps. *Geoscientific Model Development*, 17(4), 1813–1829. <https://doi.org/10.5194/gmd-17-1813-2024>
- Schiemann, R., Demory, M. E., Mizieliński, M. S., Roberts, M. J., Shaffrey, L. C., Strachan, J., & Vidale, P. L. (2014). The sensitivity of the tropical circulation and

- Maritime Continent precipitation to climate model resolution. *Climate Dynamics*, 42(9–10), 2455–2468. <https://doi.org/10.1007/s00382-013-1997-0>
- Sillmann, J., Kharin, V. V., Zhang, X., Zwiers, F. W., & Bronaugh, D. (2013). Climate extremes indices in the CMIP5 multimodel ensemble: Part 1. Model evaluation in the present climate. *Journal of Geophysical Research Atmospheres*, 118(4), 1716–1733. <https://doi.org/10.1002/jgrd.50203>
- Strandberg, G., & Lind, P. (2021). The importance of horizontal model resolution on simulated precipitation in Europe – from global to regional models. *Weather and Climate Dynamics*, 2(1), 181–204. <https://doi.org/10.5194/wcd-2-181-2021>
- Sundqvist, H. (1978). A parameterization scheme for non-convective condensation including prediction of cloud water content. *Quarterly Journal of the Royal Meteorological Society*, 104(441), 677–690. <https://doi.org/10.1002/qj.49710444110>
- Tarek, M., Brissette, F. P., & Arseneault, R. (2020). Evaluation of the ERA5 reanalysis as a potential reference dataset for hydrological modelling over North America. *Hydrology and Earth System Sciences*, 24(5), 2527–2544. <https://doi.org/10.5194/hess-24-2527-2020>
- Tetzner, D., Thomas, E., & Allen, C. (2019). A validation of ERA5 reanalysis data in the southern antarctic peninsula—Ellsworth land region, and its implications for ice core studies. *Geosciences (Switzerland)*, 9(7). <https://doi.org/10.3390/geosciences9070289>
- Torma, C., Giorgi, F., & Coppola, E. (2015). Added value of regional climate modeling over areas characterized by complex terrain-precipitation over the Alps. *Journal of Geophysical Research*, 120(9), 3957–3972. <https://doi.org/10.1002/2014JD022781>
- Wan, H., Zhang, S., Rasch, P. J., Larson, V. E., Zeng, X., & Yan, H. (2021). Quantifying and attributing time step sensitivities in present-day climate simulations conducted with EAMv1. *Geoscientific Model Development*, 14(4), 1921–1948. <https://doi.org/10.5194/gmd-14-1921-2021>
- Wang, C., Graham, R. M., Wang, K., Gerland, S., & Granskog, M. A. (2019). Comparison of ERA5 and ERA-Interim near-surface air temperature, snowfall and precipitation over Arctic sea ice: effects on sea ice thermodynamics and evolution. *Cryosphere*, 13(6), 1661–1679. <https://doi.org/10.5194/tc-13-1661-2019>
- Wehner, M. F., Reed, K. A., Li, F., Prabhat, Bacmeister, J., Chen, C. T., Paciorek, C., Gleckler, P. J., Sperber, K. R., Collins, W. D., Gettelman, A., & Jablonowski, C. (2014). The effect of horizontal resolution on simulation quality in the Community Atmospheric Model, CAM5.1. *Journal of Advances in Modeling Earth Systems*, 6(4), 980–997. <https://doi.org/10.1002/2013MS000276>
- Wehner, M. F., Smith, R. L., Bala, G., & Duffy, P. (2010). The effect of horizontal resolution on simulation of very extreme US precipitation events in a global atmosphere model. *Climate Dynamics*, 34(2), 241–247. <https://doi.org/10.1007/s00382-009-0656-y>
- Xu, X., Frey, S. K., Boluwade, A., Erler, A. R., Khader, O., Lapen, D. R., & Sudicky, E. (2019). Evaluation of variability among different precipitation products in the Northern Great Plains. *Journal of Hydrology: Regional Studies*, 24. <https://doi.org/10.1016/j.ejrh.2019.100608>
- Zhu, L., & Quiring, S. M. (2022). Exposure to precipitation from tropical cyclones has increased over the continental United States from 1948 to 2019. *Communications Earth and Environment*, 3(1). <https://doi.org/10.1038/s43247-022-00639-8>


868  
869

**Figures**



870  
871 **Fig. 1** Annual time series of the 99<sup>th</sup> percentile precipitation using observations (GPCC, green),  
872 reanalysis (ERA5, black), and model simulations (LR: blue, MR: orange, HR: red) during  
873 1982-2019 over Europe. RMSE values of 99<sup>th</sup> percentile precipitation are computed referenced  
874 to GPCC which are shown within the small bracket.  
875


876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897





Deleted:




899  
900  
901  
  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935



**Fig. 2** The 99<sup>th</sup> percentile precipitation over Europe during 1982-2019 from (a) GPCC observations, (b) ERA5 reanalysis, (c) LR, (d) MR, (e) HR, and the corresponding biases and RMSEs in (f) ERA5, (g) LR, (h) MR, and (i) HR.




938  
939



940  
941

942 **Fig. 3** RMSEs for annual total precipitation at different percentile ranges (70<sup>th</sup> – 80<sup>th</sup>, 80<sup>th</sup> –  
943 90<sup>th</sup>, 90<sup>th</sup> – 95<sup>th</sup>, 95<sup>th</sup> – 99<sup>th</sup>, 99<sup>th</sup> – 99.5<sup>th</sup>, 99.5<sup>th</sup> – 99.9<sup>th</sup> and >99.9<sup>th</sup> percentile) in ERA5 (black)  
944 and OpenIFS simulations (LR60m: magenta, LR30m: orange, LR: red, MR: green, HR: blue)  
945 referenced to GPCC during 1982-2019 over Europe. Dots are the RMSE values, and error bars  
946 are the 95 % CI.

947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968





970  
971  
972  
973

974  
975



976 **Fig. 4** Contribution of convective precipitation to extreme precipitation (>99<sup>th</sup> percentile) in (a)  
977 ERA5, (b) LR, (c) MR and (d) HR over Europe in JJA, and (e)– (g) their biases and RMSEs  
978 compared to ERA5 over the period 1982-2019.

979  
980

981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991



Deleted:



Deleted:

**Fig. 5** The same as Fig. 4 but for DJF.

994  
995

996

997  
998

999 **Fig. 6** RMSEs of total precipitation (a & d) at different percentile ranges (70<sup>th</sup> – 80<sup>th</sup>, 80<sup>th</sup> –  
1000 90<sup>th</sup>, 90<sup>th</sup> – 95<sup>th</sup>, 95<sup>th</sup> – 99<sup>th</sup>, 99<sup>th</sup> – 99.5<sup>th</sup>, 99.5<sup>th</sup> – 99.9<sup>th</sup> and >99.9<sup>th</sup>) and the corresponding  
1001 large-scale precipitation (b & e) and convective precipitation (c & f) in OpenIFS simulations  
1002 (LR60m: magenta, LR30m: orange, LR: red, MR: green, HR: blue) against ERA5 over Europe  
1003 during 1982-2019. (a) – (c) are for JJA, and (d) – (f) for DJF. Dots are the RMSE values, and  
1004 error bars are the 95 % confidence intervals. Unit is mm/d.

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014



1015

1016

1017

1018

1019



1022  
1023  
1024  
1025

**Table**

1026 Table 1: The experiment details of different horizontal resolutions and model time steps in  
1027 OpenIFS.

	LR60m	LR30m	LR	MR	HR
Vertical resolution	L91			L91	L91
Horizontal Resolution	100 km (Tco95)			50 km (Tco199)	25 km (Tco399)
Time steps	60 minutes	30 minutes	15 minutes	15 minutes	15 minutes

1028

